

Fluid Resuscitation in Sepsis

A Systematic Review and Network Meta-analysis

Bram Rochweg, MD; Waleed Alhazzani, MD; Anees Sindi, MD; Diane Heels-Ansdell, MSc; Lehana Thabane, PhD; Alison Fox-Robichaud, MD; Lawrence Mbuagbaw, MSc; Wojciech Szczeklik, MD; Fayez Alshamsi, MD; Sultan Altayyar, MD; Wang-Chun Ip, MD; Guowei Li, MSc; Michael Wang, MD; Anna Włodarczyk, MD; Qi Zhou, PhD; Gordon H. Guyatt, MD; Deborah J. Cook, MD; Roman Jaeschke, MD; and Djillali Annane, MD, PhD, for the Fluids in Sepsis and Septic Shock Group

Background: Fluid resuscitation is the cornerstone of sepsis treatment. However, whether balanced or unbalanced crystalloids or natural or synthetic colloids confer a survival advantage is unclear.

Purpose: To examine the effect of different resuscitative fluids on mortality in patients with sepsis.

Data Sources: MEDLINE, EMBASE, *ACP Journal Club*, CINAHL, HealthSTAR, the Allied and Complementary Medicine Database, and the Cochrane Central Register of Controlled Trials through March 2014.

Study Selection: Randomized trials that evaluated different resuscitative fluids in adult patients with sepsis or septic shock and death. No language restrictions were applied.

Data Extraction: Two reviewers extracted data on study characteristics, methods, and outcomes. Risk of bias for individual studies and quality of evidence were assessed.

Data Synthesis: 14 studies (18 916 patients) were included with 15 direct comparisons. Network meta-analysis at the 4-node level showed higher mortality with starches than with crystalloids (high confidence) and lower mortality with albumin than with crystalloids

(moderate confidence) or starches (moderate confidence). Network meta-analysis at the 6-node level showed lower mortality with albumin than with saline (moderate confidence) and low-molecular-weight starch (low confidence) and with balanced crystalloids than with saline (low confidence) and low- and high-molecular-weight starches (moderate confidence).

Limitations: These trials were heterogeneous in case mix, fluids evaluated, duration of fluid exposure, and risk of bias. Imprecise estimates for several comparisons in this network meta-analysis contribute to low confidence in most estimates of effect.

Conclusion: Among patients with sepsis, resuscitation with balanced crystalloids or albumin compared with other fluids seems to be associated with reduced mortality.

Primary Funding Source: The Hamilton Chapter of the Canadian Intensive Care Foundation and the Critical Care Medicine Residency Program and Critical Care Division Alternate Funding Plan at McMaster University.

Ann Intern Med. doi:10.7326/M14-0178

www.annals.org

For author affiliations, see end of text.

This article was published online first at www.annals.org on 22 July 2014.

Resuscitation with crystalloids compared with colloids for critically ill patients has been evaluated in large randomized, controlled trials (1–6) and meta-analyses (7–13). One meta-analysis (10) including 74 trials reported no difference in mortality between critically ill patients resuscitated with crystalloids and albumin (relative risk [RR], 1.01 [95% CI, 0.93 to 1.10]), hydroxyethyl starch (HES) (RR, 1.10 [CI, 0.91 to 1.32]), gelatin (RR, 0.91 [CI, 0.49 to 1.72]), or dextran (RR, 1.24 [CI, 0.94 to 1.65]). Another meta-analysis (8) reported that resuscitation with an albumin-containing solution in patients with sepsis may decrease mortality compared with solutions containing no albumin (RR, 0.82 [CI, 0.67 to 1.00]).

Recent evidence suggests that starches, compared with other fluids and regardless of molecular weight, may be associated with acute kidney injury in the general population of critically ill patients and in those with sepsis (5, 11, 13–15). A recent large pragmatic trial comparing colloids (mostly starches) with crystalloids (mostly 0.9% sodium chloride) suggested a 90-day mortality benefit with colloids (RR, 0.92 [CI, 0.86 to 0.99]) (16).

Crystalloids can be characterized on the basis of tonicity and electrolyte content. The presence of an organic anion (for example, lactate, acetate, or gluconate) and correspondingly lower chloride content that more closely resembles the composition of plasma suggest that a crystalloid is “balanced” (for example, Ringer lactate and acetate

solutions) (17). The most commonly used crystalloid, normal saline (0.9% sodium chloride), is far from “normal,” with a pH much less than 7.0 and a supraphysiologic chloride content of 154 mmol/L (18, 19). Compared with a balanced crystalloid solution, normal saline predisposes patients to hyperchloremic metabolic acidosis, decreased renal blood flow to the glomerulus, and impaired smooth-muscle contractility (20).

Investigators have not done randomized, controlled trials (RCTs) comparing balanced and unbalanced crystalloids. However, 1 large before–after study of critically ill patients showed that balanced versus unbalanced fluid solution was associated with a lower incidence of acute kidney injury (8.4% vs. 14%; $P < 0.01$) and renal replacement therapy (6.3% vs. 10%; $P = 0.05$) but no differences in hospital mortality (18).

Colloids include natural compounds, such as albumin, and synthetic compounds of HES, gelatin, or dextran. Expansion of plasma volume increases in proportion to the

See also:

Editorial comment. 1

Web-Only
Supplements

osmotic or oncotic potential, and colloids theoretically require less volume than crystalloids to achieve equivalent hemodynamic effect (19). Limitations of colloids include development of acute kidney injury and coagulation disorders with starches (14) and the risk for exposure to blood products with albumin (19). Another important consideration is the biochemical properties of the crystalloid solution in which the colloid is dissolved. For example, the chloride concentrations in HES may vary between 154 mmol/L (Voluven, Fresenius Kabi) and 118 mmol/L (Tetraspan, B. Braun Medical) (21).

Whether any of these fluid properties translate into a survival advantage remains unclear, particularly regarding the optimal fluid for resuscitation in patients with sepsis. Fluid resuscitation, in addition to antibiotics and source control, is a cornerstone of initial management of sepsis (22). However, fluid management in patients with sepsis varies widely in practice (16, 23, 24).

Meta-analyses of fluid resuscitation have been limited by not focusing on patients with sepsis (7, 9, 10), not considering electrolyte composition (5, 8, 10, 11), considering only 2 or 3 categories of fluid (25), not including direct and indirect comparisons in the same model, and omission of recent large RCTs (3–5, 16). Therefore, we did a network meta-analysis (NMA) considering direct and indirect comparisons of all types of fluid resuscitation tested in RCTs in patients with severe sepsis and septic shock, focusing on the effect of these interventions on mortality.

METHODS

Data Sources and Searches

This review was done using a predefined protocol. Initially, we searched MEDLINE (1948 to December 2012), EMBASE (1980 to December 2012), *ACP Journal Club* (1991 to December 2012), the Cochrane Central Register of Controlled Trials, HealthSTAR, the Allied and Complementary Medicine Database, and CINAHL. We updated the MEDLINE and EMBASE searches in August 2013 and March 2014. We screened previously published meta-analyses for relevant citations. **Supplement 1** (available at www.annals.org) presents the search terms used.

Six reviewers working in 3 pairs screened the titles and abstracts to determine potential eligibility, and entries identified by any reviewer proceeded to the full-text eligibility review. Pretested eligibility forms were used for full-text review, which was also done in duplicate. A third adjudicator helped to resolve disagreements through consensus.

Study Selection

We selected parallel-group RCTs, including factorial designs, but excluded quasi-randomized and crossover trials. We excluded all studies published by Dr. Joachim Boldt because of suspected lack of integrity (26, 27). We

did not apply restrictions on language or publication date.

We included studies that involved adult (aged ≥ 16 years) critically ill patients with severe sepsis or septic shock as defined by the investigators and who required fluid resuscitation (defined as the administration of a bolus of intravenous fluid exceeding the amount required for maintenance or replacement fluids). We included studies with mixed critically ill populations whenever separate data for patients with sepsis were available. We excluded studies in which most patients had the systemic inflammatory response syndrome secondary to other causes (such as burn, pancreatitis, and trauma) without a clear sepsis subgroup and those focusing on patients after elective surgery.

Interventions studied included any fluid or fluid strategy used for resuscitation compared with another fluid or fluid strategy. We excluded studies in which the primary goal was to assess short-term hemodynamic response.

Our outcome was 90-day mortality or, if not available, 30-day, intensive care unit, or hospital mortality, whichever was longest.

Data Extraction

Pairs from the same 6 reviewers abstracted data in duplicate. Another clinician reviewed disagreements, and consensus was reached by discussion. We contacted authors of primary publications for missing or unclear information.

Risk of Bias

Independently and in duplicate, reviewers assessed risk of bias using a modified version of the Cochrane Collaboration assessment tool (28, 29). We judged each included study as having low, probably low, probably high, or high risk of bias for randomization-sequence generation, randomization concealment, blinding, incomplete data, selective reporting, and free of other bias (including intention-to-treat analysis). The overall rating of risk of bias for each study was the lowest rating for any of the criteria (**Appendix Table**, available at www.annals.org).

Data Synthesis and Analysis

Our analysis classified fluids as crystalloids (divided into balanced and unbalanced solutions) and colloids (divided into albumin, gelatin, and low- and high-molecular-weight HES [threshold molecular weight, 150 000 kDa]). We considered fluid balanced if it contained an anion of a weak acid (buffer) and its chloride content was correspondingly less than in 0.9% sodium chloride (21). The relevant analyses were a 4-node NMA (crystalloids vs. albumin vs. HES vs. gelatin), a 6-node NMA (crystalloids vs. albumin vs. HES vs. gelatin, with crystalloids divided into balanced or unbalanced and HES divided into low or high molecular weight), and a conventional direct frequentist fixed-effects meta-analytic comparison of crystalloids versus colloids.

To calculate direct estimates of treatment effect for each pair of treatments within the 4- and 6-node networks,

we did a frequentist fixed-effects meta-analysis. We reported the results as odds ratios (ORs) and corresponding 95% CIs. We evaluated heterogeneity by estimating the variance between studies (chi-square test and I^2 statistic) (30, 31).

Using a Bayesian framework, we did 4- and 6-node fixed-effects NMAs for each treatment. We reported the results as ORs and corresponding 95% credibility intervals (CrIs), which are the Bayesian analogue of the 95% CIs (32). The ORs reported are relative effects of compared fluids. The models are based on 80 000 iterations with a burn-in of 40 000 and a thin of 10. We used a random seed and vague priors. We assessed nonconvergence on the basis of Brooks–Gelman–Rubin plots (33).

We used the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) approach to assess confidence in estimates of effect (quality of evidence) associated with specific comparisons, including estimates from direct, indirect, and final NMAs (Supplement 2, available at www.annals.org) (34). Our confidence assessment addressed risk of bias, incoherence, imprecision, inconsistency, indirectness, and publication bias.


The starting point for confidence in direct and indirect estimates was “high.” However, indirect estimates were potentially rated down for intransitivity (that is, differences in patients, co-interventions, or setting that could lead to effect modification and thus a misleading comparison of fluid management strategies). We inferred confidence in indirect estimates by examining the connecting loops associated with the particular comparison.

The confidence rating chosen was the lowest of the direct estimates contributing to the indirect comparison. For example, consider a comparison of A versus B that is informed by comparisons of A versus C and B versus C. If A versus C was rated as high confidence and B versus C as moderate confidence, the overall indirect confidence rating was initially based on the comparison of B versus C and was then potentially rated down to low for indirectness. Precision was judged on the basis of the CrI around the point estimate from the indirect comparison.

The overall NMA confidence rating was the higher confidence in the direct and indirect comparisons with the possibility of rating up further for gains in precision with pooling of direct and indirect comparisons. Confidence was lowered 1 level for incoherence if the CrIs between the direct and indirect estimates did not significantly overlap. The approach we used was consistent with preliminary GRADE guidance and GRADE methods for direct comparisons (35, 36).

We did the Bayesian analysis for the NMA using WinBUGS 1.4.3 (Medical Research Council Biostatistics Unit; www.mrc-bsu.cam.ac.uk/software/bugs/), and all frequentist analysis was done using Review Manager 5.2.3 (Nordic Cochrane Centre; <http://ims.cochrane.org/revman/download>).

Figure. Summary of evidence search and selection.



AMED = Allied and Complementary Medicine Database; CENTRAL = Cochrane Central Register of Controlled Trials; RCT = randomized, controlled trial.

Role of the Funding Source

This research was funded by the Hamilton Chapter of the Canadian Intensive Care Foundation and the Critical Care Medicine Residency Program and Critical Care Division Alternate Funding Plan, both at McMaster University. The funding sources had no role in the design, conduct, or reporting of this study or in the decision to submit the manuscript for publication.

RESULTS

Of 9875 titles identified during the primary search (7329 after duplicates were removed), 7146 were judged as ineligible, leaving 185 studies for full-text review (Figure).

Table 1. Study Characteristics

Study, Year (Reference)	Centers, <i>n</i>	Country	Randomly Assigned Patients, <i>n</i>	Sepsis*	Mean APACHE II Score
Haupt et al, 1982 (38)	1	United States	17	NR	NR
Rackow et al, 1989 (42)	1	United States	20	Yes	NR
Schortgen et al, 2001 (43)	3	France	129	Yes	NR
Finfer et al, 2004 (2)	16	Australia and New Zealand	6997	Yes	18.9
Brunkhorst et al, 2008 (1)	18	Germany	537	Yes	20.0
Li et al, 2008 (39)	1	China	60	Yes	18.0
McIntyre et al, 2008 (41)	4	New Zealand and Canada	40	Yes	20.6
Dubin et al, 2010 (37)	2	Argentina	25	Yes	NR**
Siegemund, 2011 (44)	1	Switzerland	241	NR	Not included
Guidet et al, 2012 (6)	24	France and Germany	196	Yes	NR**
Lv et al, 2012 (40)	1	China	42	NR	NR
Myburgh et al, 2012 (3)	32	Australia and New Zealand	7000	NR	17.0
Perner et al, 2012 (4)	26	Scandinavia	804	Yes	NR§§
Anname et al, 2013 (16)	57	Worldwide	2857	Yes	NR

AL = albumin; APACHE = Acute Physiology and Chronic Health Evaluation; COL = colloid; CRY = crystalloid; GL = gelatin; HES = hydroxyethyl starch; H-HES = high-molecular-weight HES; HS = hypertonic saline; ICU = intensive care unit; L-HES = low-molecular-weight HES; NR = not reported; NS = normal saline; RA = Ringer acetate; RL = Ringer lactate.

* According to international sepsis definition, or similar. Definition from reference 45.

† Except where noted.

‡ Mean.

§ Hospital stay.

|| ICU stay.

¶ The intervention in this study was unblinded.

** Mean Sequential Organ Failure Assessment score, 8.5.

†† Median.

‡‡ Study fluid administered during the first 4 d of the study.

§§ Mean Sequential Organ Failure Assessment score, 7.0.

||| Mean Sequential Organ Failure Assessment score, 8.0.

This total included 2 unpublished studies, 1 found through screening titles included in a previous meta-analysis (5) and another (since published) found through author correspondence (16). Of these 185 RCTs, 171 were ineligible, leaving 14 eligible RCTs (1–6, 16, 37–43).

Table 1 summarizes the characteristics of 14 trials involving 18 916 adults available for analysis (44). In 4 included trials (2, 3, 16, 38), septic patients were a subgroup of all patients enrolled; only 1 of these 4 studies (16) stratified randomization on the basis of the diagnosis of sepsis. Through contact with primary authors or authors of previous reviews, we obtained relevant data for 10 of the 14 studies. No additional data were needed for the other 4 studies.

Most included studies used a definition of sepsis consistent with international consensus (45). Five studies reported mean Acute Physiology and Chronic Health Eval-

uation II scores between 17.0 and 20.6, whereas another 4 studies reported median Sequential Organ Failure Assessment scores between 7.0 and 8.5. The fluid intervention period lasted between 24 hours and 90 days. The observation period for outcome assessment varied between 30 days and 1 year.

Appendix Figures 1 and 2 (available at www.annals.org) present network nodes showing direct and indirect comparisons and the number of studies available for comparisons. Table 2 indicates the contribution of individual studies and the subclassification of each fluid at each level of analysis.

4-Node Analysis

Table 3 presents the results of the 4-node analysis. The results suggest higher mortality with starches than with crystalloids (OR, 1.13 [95% CrI, 0.99 to 1.30]; high

Table 1—Continued

Intervention and Cumulative Dose	Blood Products Transfused, n/N (%)†	Intervention Period (Mortality Observation Period)	Overall Risk of Bias	Deaths, n/N	Industry Sponsor
NS: 6371 mL‡ AL: 3134 mL‡ H-HES: 4466 mL‡	NR	24 h§	Probably low	NS: 3/4 AL: 5/7 H-HES: 3/6	Cutter
AL: 975 mL H-HES: 900 mL	NR	<24 h§	Probably low	AL: 5/10 H-HES: 5/10	Dupont Critical Care; Cutter
GL: 43 mL/kg H-HES: 31 mL/kg	NR	GL: Until ICU discharge H-HES: 4 d	Low	GL: 29/64 H-HES: 28/65	NR
NS: >3000 mL AL: >2000 mL	NR	28 d (31 d)	Low	NS: 217/615 AL: 185/603	CSL Behring
RL: 1.32 × HES dose HES: 70.4 mL/kg	RL: 189/275 (68.7) HES: 199/262 (76)	21 d (90 d)	Low¶	RL: 93/274 HES: 107/261	B. Braun Medical; Novo Nordisk; HemoCue
HES/HS: NR HES: NR HS: NR NS: NR	NR	<24 h (31 d)	Probably low¶	HES/HS: 5/15 HES: 9/15 HS: 10/15 NS: 10/15	NR
NS: 2100 mL H-HES: 1900 mL	NS: 5/19 (26) H-HES: 10/21 (48)	<24 h§	Low	NS: 7/19 H-HES: 9/21	Bristol-Myers Squibb; Edward Life Sciences
NS: 6254 mL L-HES: 2610 mL	NS: 18% L-HES: 22%	24 h (31 d)	Probably low¶	NS: 7/13 L-HES: 3/12	NR
NS: NR L-HES: 3775 mL††	NR	5 d (1 y)	Low	NS: 50/124 L-HES: 44/117	Fresenius
NS: 2788 mL L-HES: 2615 mL	NS: 20/96 (21) L-HES: 29/100 (29)	4 d (90 d)	Low	NS: 32/95 L-HES: 40/99	Fresenius Kabi
RL: 3460 mL L-HES: 2770 mL	NR	Not clear	Probably low¶	RL: 12/20 L-HES: 7/22	NR
NS: 2456 mL‡‡ L-HES: 2104 mL‡‡	NR	90 d (90 d)	Low	NS: 224/945 L-HES: 248/976	Fresenius Kabi
RA: 3000 mL† L-HES: 3000 mL†	RA: 204/380 (54) L-HES: 243/376 (65)	90 d (90 d)	Low	RA: 173/400 L-HES: 202/398	B. Braun Medical
CRY: 3000 mL in 7 d COL: 2000 mL in 7 d	CRY: 358/1443 (24.8) COL: 377/1414 (26.7)	Until ICU discharge (90 d)	Low	CRY: 286/779 COL: 252/774	None

confidence) and lower mortality with albumin than with crystalloids (OR, 0.83 [CrI, 0.65 to 1.04]; moderate confidence) and starches (OR, 0.73 [CrI, 0.56 to 0.95]; moderate confidence) (Table 3).

6-Node Analysis

Table 4 presents the results of the 6-node analysis. Evidence suggests that albumin is superior to saline (OR, 0.82 [CrI, 0.65 to 1.04]; moderate confidence) and low-molecular-weight starch (OR, 0.79 [CrI, 0.59 to 1.06]; low confidence). It also suggests that balanced crystalloids are superior to saline (OR, 0.78 [CrI, 0.58 to 1.05]; low confidence), high-molecular-weight starch (OR, 0.82 [CrI, 0.60 to 1.13]; moderate confidence), and low-molecular-weight starch (OR, 0.75 [CrI, 0.58 to 0.97]; moderate confidence).

Post Hoc Sensitivity Analysis

Supplement 3 (available at www.annals.org) shows the results of 2 sensitivity analyses that were done at the request of peer reviewers. The first analysis incorporates the recently published ALBIOS (Albumin Italian Outcome Sepsis) trial (46). Given that the aim of fluid administration in this trial was not clearly resuscitative, we had initially excluded it from our review. Inclusion of the ALBIOS trial in this post hoc analysis did not affect our results. The second analysis was done to investigate the

effect of excluding the older studies by Haupt and Rackow (38) and Rackow and colleagues (42), both with low event rates and few patients. This requested post hoc analysis also did not affect our results.

Crystalloids Versus Colloids

A crude analysis showed no difference in mortality between colloids and crystalloids for fluid resuscitation in adult patients (OR, 0.99 [CI, 0.89 to 1.10]; $P = 0.85$; chi-square = 23.20; $I^2 = 53%$; moderate confidence [because of inconsistency]) (Appendix Figure 3, available at www.annals.org).

DISCUSSION

The results of this NMA highlight potentially important differences in mortality between crystalloid solutions. Our findings suggest an advantage of balanced crystalloids versus saline (low confidence) and low- or high-molecular-weight starch (moderate confidence), with similar mortality results for balanced crystalloids and albumin (very low confidence) (Table 4). These differences were not detectable using a standard meta-analytic approach directly comparing “any crystalloids versus any colloids,” the results of which are consistent with previous meta-analyses (7, 10) and a recent large RCT (16).

Table 2. Contribution of Individual Studies to Each Level of Analysis

Fluid Name, by Study, Year (Reference)	Trade Name	Crystalloid vs. Colloid	4-Node Analysis	6-Node Analysis
Haupt and Rackow, 1982 (38)				
Normal saline	Saline	Crystalloid	Crystalloid	Saline
5% albumin	Albumin	Colloid	Albumin	Albumin
6% HES	Hespan (B. Braun Medical)	Colloid	HES	H-HES
Rackow et al, 1989 (42)				
5% albumin	Albumin	Colloid	Albumin	Albumin
10% HES	Pentastarch	Colloid	HES	H-HES
Schortgen et al, 2001 (43)				
3% gelatin	Plasmagel (Laboratoire Roger Bellon)	Colloid	Gelatin	Gelatin
6% HES	HAES-steril (Fresenius Kabi)	Colloid	HES	H-HES
Finfer et al, 2004 (2)				
Normal saline	Saline	Crystalloid	Crystalloid	Saline
4% albumin	Albumex (CSL Behring)	Colloid	Albumin	Albumin
Brunkhorst et al, 2008 (1)				
RL	RL	Crystalloid	Crystalloid	Balanced crystalloid
10% HES	Pentastarch	Colloid	HES	H-HES
Li et al, 2008 (39)				
HES in hypertonic saline	NR	Colloid	HES	H-HES
HES	NR	Colloid	HES	H-HES
Hypertonic saline	NR	Crystalloid	Crystalloid	Saline
Normal saline	NR	Crystalloid	Crystalloid	Saline
McIntyre et al, 2008 (41)				
Normal saline	Saline	Crystalloid	Crystalloid	Saline
10% HES	Pentastarch	Colloid	HES	H-HES
Dubin et al, 2010 (37)				
Normal saline	Saline	Crystalloid	Crystalloid	Saline
6% HES	Voluven (Fresenius Kabi)	Colloid	HES	L-HES
Siegemund, 2011 (44)				
Normal saline	Saline	Crystalloid	Crystalloid	Saline
6% HES	Voluven (Fresenius Kabi)	Colloid	HES	L-HES
Guidet et al, 2012 (6)				
Normal saline	Saline	Crystalloid	Crystalloid	Saline
6% HES	Voluven (Fresenius Kabi)	Colloid	HES	L-HES
Lv et al, 2012 (40)				
RL	RL	Crystalloid	Crystalloid	Balanced crystalloid
6% HES	Voluven (Fresenius Kabi)	Colloid	HES	L-HES
Myburgh et al, 2012 (3)				
Normal saline	Saline	Crystalloid	Crystalloid	Saline
6% HES	Voluven (Fresenius Kabi)	Colloid	HES	L-HES
Perner et al, 2012 (4)				
RA	RA	Crystalloid	Crystalloid	Balanced crystalloid
6% HES	Tetraspan (B. Braun Medical)	Colloid	HES	L-HES
Annane et al, 2013 (16)				
Any crystalloid	NA	Crystalloid	Not included in analysis	Not included in analysis
Any colloid	NA	Colloid	Not included in analysis	Not included in analysis

HES = hydroxyethyl starch; H-HES = high-molecular-weight HES; L-HES = low-molecular-weight HES; NA = not applicable; RL = Ringer lactate; NR = not reported.

Biological rationale is consistent with the findings of lower mortality with balanced crystalloid solutions than with saline, because balanced fluids mimic the homeostatic composition of body fluids to a greater extent than unbalanced fluids (17, 18). Which component of balanced solutions—the pH, electrolyte composition, or presence of a buffer—contributes to this potential benefit is unclear.

These results raise concerns about whether using mostly unbalanced crystalloids in the acute resuscitation of patients with sepsis is optimal. Our findings may partially explain the higher 90-day mortality rates observed in the crystalloid group compared with the colloid group of a recent trial in which 86% of the patients randomly assigned to crystalloids received normal saline (16).

Table 3. NMA Results of 4-Node Analysis, Including Confidence Assessments

Comparison	Trials With Direct Comparisons, <i>n</i>	Direct Estimate (95% CI); Quality of Evidence	Indirect Estimate (95% CrI); Quality of Evidence	NMA Estimate (95% CrI)*; Quality of Evidence
Starch vs. crystalloid	10	1.14 (0.99–1.30); High	0.81 (0.13–5.14); Very low†‡	1.13 (0.99–1.30); High
Albumin vs. crystalloid	2	0.81 (0.64–1.03); Moderate§	1.13 (0.18–7.32); Very low†‡	0.83 (0.65–1.04); Moderate
Gelatin vs. crystalloid	0	–	1.24 (0.61–2.55); Very low†§	1.24 (0.61–2.55); Very low
Albumin vs. starch	2	1.40 (0.35–5.56); Low‡	0.71 (0.54–0.94); Moderate†	0.73 (0.56–0.95); Moderate
Gelatin vs. starch	1	1.09 (0.55–2.19); Low‡	–	1.10 (0.54–2.22); Low
Gelatin vs. albumin	0	–	1.51 (0.71–3.20); Very low†‡	1.51 (0.71–3.20); Very low

CrI = credibility interval; NMA = network meta-analysis.

* Higher of direct or indirect confidence.

† Rated down for indirectness.

‡ Rated down 2 levels for imprecision.

§ Rated down for imprecision.

We found that the effects of different colloids may also vary, with albumin seeming equivalent or superior to all alternatives. Starches, regardless of molecular weight, seem inferior to alternative resuscitation fluids (that is, albumin and balanced crystalloids). Data on gelatin are markedly less robust. Gelatin is associated with increased mortality relative to some other resuscitation fluids; however, only 1 small trial contributes directly to this analysis. The heterogeneity of effects observed across different types of colloids renders crude comparisons of colloids versus crystalloids uninformative (Appendix Figure 3).

The strengths of this review include a precise clinical question restricted to patients with sepsis rather than all critically ill patients. It focuses on resuscitation rather than maintenance fluid and distinguishes between balanced and unbalanced crystalloids and between starches of different molecular weight. We did a comprehensive search and risk-of-bias assessment, with both processes involving duplicate review and third-party adjudication. Using rigorous NMA methods (32), we incorporated direct and indirect

evidence. The GRADE approach, seldom applied to NMA, allowed reporting of confidence in estimates of effect when interpreting each unique fluid comparison.

The limitations of our review include the small number of studies relative to the number of comparisons considered, resulting in low confidence in estimates for many key analyses. Although all included studies focused on fluid for resuscitation, fluid protocols were somewhat heterogeneous, with the amounts of fluid administered and durations of the fluid intervention varying.

Some observed results may be related to the interplay of different fluid properties, particularly the differential presence of chloride in each colloid. For example, the albumin used in the largest albumin study (2) was dissolved in a crystalloid solution containing greater than 6 mmol of caprylate and 128 mmol of chloride per liter and suggested a trend toward benefit compared with saline (chloride content, 154 mmol/L of chloride) in the sepsis subgroup (47). Given our results suggesting lower mortality associated with balanced solutions, the apparent trend favoring albu-

Table 4. NMA Results of 6-Node Analysis, Including Confidence Assessments

Comparison	Trials With Direct Comparisons, <i>n</i>	Direct Estimate (95% CI); Quality of Evidence	Indirect Estimate (95% CrI); Quality of Evidence	NMA Estimate (95% CrI)*; Quality of Evidence
L-HES vs. saline	4	1.07 (0.89–1.29); Moderate†	0.59 (0.25–1.35); Very low†‡§	1.04 (0.87–1.25); Moderate
H-HES vs. saline	3	0.64 (0.30–1.37); Moderate†	1.13 (0.71–1.80); Very low†‡	0.95 (0.64–1.41); Moderate
Albumin vs. saline	2	0.81 (0.64–1.03); Moderate†	0.96 (0.14–6.31); Very low†	0.82 (0.65–1.04); Moderate
Balanced crystalloid vs. saline	0	–	0.78 (0.58–1.05); Low†‡	0.78 (0.58–1.05); Low
Gelatin vs. saline	0	–	1.04 (0.46–2.32); Very low†‡	1.04 (0.46–2.32); Very low
H-HES vs. L-HES	0	–	0.91 (0.63–1.33); Low†‡	0.91 (0.63–1.33); Low
Albumin vs. L-HES	0	–	0.79 (0.59–1.06); Low†‡	0.79 (0.59–1.06); Low
Balanced crystalloid vs. L-HES	2	0.80 (0.61–1.04); Moderate§	0.44 (0.19–0.97); Moderate‡	0.75 (0.58–0.97); Moderate
Gelatin vs. L-HES	0	–	1.00 (0.44–2.21); Very low†‡	1.00 (0.44–2.21); Very low
Albumin vs. H-HES	2	1.40 (0.35–5.56); Low	0.83 (0.52–1.33); Low†‡	0.87 (0.55–1.36); Low
Balanced crystalloid vs. H-HES	1	0.74 (0.52–1.05); Moderate†	1.35 (0.63–2.92); Very low†	0.82 (0.60–1.13); Moderate
Gelatin vs. H-HES	1	1.09 (0.55–2.19); Low	–	1.10 (0.54–2.21); Low
Balanced crystalloid vs. albumin	0	–	0.95 (0.65–1.38); Very low†‡	0.95 (0.65–1.38); Very low
Gelatin vs. albumin	0	–	1.26 (0.55–2.90); Very low†	1.26 (0.55–2.90); Very low
Gelatin vs. balanced crystalloid	0	–	1.34 (0.61–2.89); Very low†	1.34 (0.61–2.89); Very low

CrI = credibility interval; H-HES = high-molecular-weight hydroxyethyl starch; L-HES = low-molecular-weight hydroxyethyl starch; NMA = network meta-analysis.

* Higher of direct or indirect confidence.

† Rated down for imprecision.

‡ Rated down for indirectness.

§ Rated down for inconsistency ($I^2 = 80\%$; $P = 0.03$ for heterogeneity).

|| Rated down 2 levels for imprecision.

min may be at least partly related to the solution in which it was dissolved. In a similar manner, 1 study on starches (1) found statistically significant benefit of balanced crystalloid that contained 45 mmol of lactate and 106 mmol of chloride per liter compared with starch dissolved in saline (chloride content, 154 mmol/L). The mortality difference may be partly due to the balanced versus unbalanced nature of the solutions.

This analysis provides a current, comprehensive summary of the effect of resuscitation fluids on mortality in patients with sepsis. The presence of buffering substances and chloride content is often overlooked when choosing resuscitative fluids in the clinical setting and is rarely transparently reported in clinical trials, which should no longer be the case. Our analyses suggest that balanced solutions may be preferable to unbalanced solutions if crystalloids are used and that, in sepsis, albumin may be a reasonable alternative to other resuscitation fluids. However, relative to balanced crystalloids, albumin confers a small risk associated with transfusion of blood products and costs markedly more.

This NMA suggests that clinicians should be aware of the possible effect of the mineral content and the presence or absence of buffering anions in resuscitation fluids. Future trials on resuscitation for septic shock should evaluate the pH and chloride content of the fluids being compared and are needed to confirm or refute these findings.

From McMaster University, Hamilton, Ontario, Canada; Prince Sultan Military Medical City, Riyadh, Saudi Arabia; King Abdulaziz University, Jeddah, Saudi Arabia; Jagiellonian University Medical College, Krakow, Poland; and University of Versailles Saint-Quentin-en-Yvelines, Garches, France.

Note: Dr. Cook coauthored a fluid trial cited in this review, and Dr. Annane was principal investigator of a fluid trial cited in this review.

Acknowledgment: The authors thank Drs. J.L. Vincent (38, 42), L. Jie (40), S. Finfer (2), K. Reinhart (1), A. Chopra and J.L. Falk (38, 42), F. Schortgen (43), B. Wills and N. Haase (5, 6), L.L. McIntyre (41), and K. Maitland and J. Myburgh (3) for providing information contributing to this article. They also thank librarians Lois Cottrell and Jean Maragno for their invaluable help with structuring and performing their search.

Financial Support: By the Hamilton Chapter of the Canadian Intensive Care Foundation and the Critical Care Medicine Residency Program and Critical Care Division Alternate Funding Plan, both at McMaster University. Dr. Cook is supported by the Canadian Institutes of Health Research.

Disclosures: Disclosures can be viewed at www.acponline.org/authors/icmje/ConflictOfInterestForms.do?msNum=M14-0178.

Requests for Single Reprints: Deborah J. Cook, MD, Academic Chair, Critical Care Medicine, Room D176, St. Joseph's Healthcare, 50 Charlton Avenue East, Hamilton, Ontario L8N 3A5, Canada; e-mail, debcook@mcmaster.ca.

Current author addresses and author contributions are available at www.annals.org.

References

1. Brunkhorst FM, Engel C, Bloos F, Meier-Hellmann A, Ragaller M, Weiler N, et al; German Competence Network Sepsis (SepNet). Intensive insulin therapy and pentastarch resuscitation in severe sepsis. *N Engl J Med*. 2008;358:125-39. [PMID: 18184958] doi:10.1056/NEJMoa070716
2. Finfer S, Bellomo R, Boyce N, French J, Myburgh J, Norton R; SAFE Study Investigators. A comparison of albumin and saline for fluid resuscitation in the intensive care unit. *N Engl J Med*. 2004;350:2247-56. [PMID: 15163774]
3. Myburgh JA, Finfer S, Bellomo R, Billot L, Cass A, Gattas D, et al; CHEST Investigators. Hydroxyethyl starch or saline for fluid resuscitation in intensive care. *N Engl J Med*. 2012;367:1901-11. [PMID: 23075127] doi:10.1056/NEJMoa1209759
4. Perner A, Haase N, Guttormsen AB, Tenhunen J, Klemenzson G, Åneman A, et al; 6S Trial Group. Hydroxyethyl starch 130/0.42 versus Ringer's acetate in severe sepsis. *N Engl J Med*. 2012;367:124-34. [PMID: 22738085]
5. Haase N, Perner A, Hennings LI, Siegemund M, Lauridsen B, Wetterslev M, et al. Hydroxyethyl starch 130/0.38-0.45 versus crystalloid or albumin in patients with sepsis: systematic review with meta-analysis and trial sequential analysis. *BMJ*. 2013;346:f839. [PMID: 23418281] doi:10.1136/bmj.f839
6. Guidet B, Martinet O, Boulain T, Philippart F, Poussel JF, Maizel J, et al. Assessment of hemodynamic efficacy and safety of 6% hydroxyethylstarch 130/0.4 vs. 0.9% NaCl fluid replacement in patients with severe sepsis: The CRYSTMAS study. *Crit Care*. 2012;16:R94. [PMID: 22624531]
7. Bunn F, Trivedi D. Colloid solutions for fluid resuscitation. *Cochrane Database Syst Rev*. 2012;7:CD001319. [PMID: 22786474] doi:10.1002/14651858.CD001319.pub5
8. Delaney AP, Dan A, McCaffrey J, Finfer S. The role of albumin as a resuscitation fluid for patients with sepsis: a systematic review and meta-analysis. *Crit Care Med*. 2011;39:386-91. [PMID: 21248514] doi:10.1097/CCM.0b013e3181ffe217
9. Gattas DJ, Dan A, Myburgh J, Billot L, Lo S, Finfer S; CHEST Management Committee. Fluid resuscitation with 6% hydroxyethyl starch (130/0.4) in acutely ill patients: an updated systematic review and meta-analysis. *Anesth Analg*. 2012;114:159-69. [PMID: 22184610] doi:10.1213/ANE.0b013e318236b4d6
10. Perel P, Roberts I, Ker K. Colloids versus crystalloids for fluid resuscitation in critically ill patients. *Cochrane Database Syst Rev*. 2013;2:CD000567. [PMID: 23450531] doi:10.1002/14651858.CD000567.pub6
11. Zarychanski R, Abou-Setta AM, Turgeon AF, Houston BL, McIntyre L, Marshall JC, et al. Association of hydroxyethyl starch administration with mortality and acute kidney injury in critically ill patients requiring volume resuscitation: a systematic review and meta-analysis. *JAMA*. 2013;309:678-88. [PMID: 23423413] doi:10.1001/jama.2013.430
12. Wilkes MM, Navickis RJ. Patient survival after human albumin administration. A meta-analysis of randomized, controlled trials. *Ann Intern Med*. 2001;135:149-64. [PMID: 11487482]
13. Bellomo R, Prowle JR, Echeverri JE, Ligabo V, Ronco C. Fluid management in septic acute kidney injury and cardiorenal syndromes. *Contrib Nephrol*. 2010;165:206-18. [PMID: 20427971] doi:10.1159/000313760
14. Haase N, Perner A. Hydroxyethyl starch for resuscitation. *Curr Opin Crit Care*. 2013;19:321-5. [PMID: 23743591] doi:10.1097/MCC.0b013e3283632de6
15. U.S. Food and Drug Administration. FDA Safety Communication: boxed warning on increased mortality and severe renal injury, and additional warning on risk of bleeding, for use of hydroxyethyl starch solutions in some settings. 25 November 2013. Accessed at www.fda.gov/biologicsbloodvaccines/safetyavailability/ucm358271.htm on 6 September 2013.
16. Annane D, Siami S, Jaber S, Martin C, Elatrous S, Declère AD, et al; CRISTAL Investigators. Effects of fluid resuscitation with colloids vs crystalloids on mortality in critically ill patients presenting with hypovolemic shock: the CRISTAL randomized trial. *JAMA*. 2013;310:1809-17. [PMID: 24108515] doi:10.1001/jama.2013.280502
17. Morgan TJ. The ideal crystalloid—what is 'balanced'? *Curr Opin Crit Care*. 2013;19:299-307. [PMID: 23743589] doi:10.1097/MCC.0b013e3283632d46

18. Yunos NM, Bellomo R, Hegarty C, Story D, Ho L, Bailey M. Association between a chloride-liberal vs chloride-restrictive intravenous fluid administration strategy and kidney injury in critically ill adults. *JAMA*. 2012;308:1566-72. [PMID: 23073953] doi:10.1001/jama.2012.13356
19. Myburgh JA, Mythen MG. Resuscitation fluids. *N Engl J Med*. 2013;369:1243-51. [PMID: 24066745] doi:10.1056/NEJMr1208627
20. Lobo DN. Intravenous 0.9% saline and general surgical patients: a problem, not a solution [Editorial]. *Ann Surg*. 2012;255:830-2. [PMID: 22470080] doi:10.1097/SLA.0b013e318250766c
21. Rochweg B, Wludarczyk A, Szczeklik W, Alhazzani W, Sindi A, Alshamsi F, et al; FISSH group. Fluid resuscitation in severe sepsis and septic shock: systematic description of fluids used in randomized trials. *Pol Arch Med Wewn*. 2013;123:603-8. [PMID: 24185099]
22. Rivers E, Nguyen B, Havstad S, Ressler J, Muzzin A, Knoblich B, et al; Early Goal-Directed Therapy Collaborative Group. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med*. 2001;345:1368-77. [PMID: 11794169]
23. Jiwaji Z, Brady S, McIntyre LA, Gray A, Walsh TS. Emergency department management of early sepsis: a national survey of emergency medicine and intensive care consultants. *Emerg Med J*. 2013. [PMID: 24005642] doi:10.1136/emered-2013-202883
24. Jones D, McEvoy S, Merz TM, Higgins A, Bellomo R, Cooper JD, et al. International albumin use: 1995 to 2006. *Anaesth Intensive Care*. 2010;38:266-73. [PMID: 20369758]
25. Bansal M, Farrugia A, Balboni S, Martin G. Relative survival benefit and morbidity with fluids in severe sepsis—a network meta-analysis of alternative therapies. *Curr Drug Saf*. 2013;8:236-45. [PMID: 23909705]
26. Miller DR. Update to readers and authors on ethical and scientific misconduct: retraction of the “Boldt articles” [Editorial]. *Can J Anaesth*. 2011;58:777-81. [PMID: 21800211] doi:10.1007/s12630-011-9558-7
27. Reilly C. Retraction. Notice of formal retraction of articles by Dr. Joachim Boldt. *Br J Anaesth*. 2011;107:116-7. [PMID: 21685126] doi:10.1093/bja/aer068
28. Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al; Cochrane Bias Methods Group. The Cochrane Collaboration’s tool for assessing risk of bias in randomised trials. *BMJ*. 2011;343:d5928. [PMID: 22008217] doi:10.1136/bmj.d5928
29. Guyatt GH, Busse J. Methods commentary: risk of bias in randomized trials 1. Evidence Partners. 2011. Accessed at <http://distillercer.com/resources/methodological-resources/risk-of-bias-commentary/> on 13 September 2013.
30. Jansen JP, Cope S. Meta-regression models to address heterogeneity and inconsistency in network meta-analysis of survival outcomes. *BMC Med Res Methodol*. 2012;12:152. [PMID: 23043545] doi:10.1186/1471-2288-12-152
31. Jansen JP, Naci H. Is network meta-analysis as valid as standard pairwise meta-analysis? It all depends on the distribution of effect modifiers. *BMC Med*. 2013;11:159. [PMID: 23826681] doi:10.1186/1741-7015-11-159
32. Mills EJ, Ioannidis JP, Thorlund K, Schünemann HJ, Puhan MA, Guyatt GH. How to use an article reporting a multiple treatment comparison meta-analysis. *JAMA*. 2012;308:1246-53. [PMID: 23011714]
33. Gelman A, Rubin DB. Inferences from iterative simulation using multiple sequences. *Stat Sci*. 1992;7:457-72.
34. Balshem H, Helfand M, Schünemann HJ, Oxman AD, Kunz R, Brozek J, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol*. 2011;64:401-6. [PMID: 21208779] doi:10.1016/j.jclinepi.2010.07.015
35. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al; GRADE Working Group. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336:924-6. [PMID: 18436948] doi:10.1136/bmj.39489.470347.AD
36. Guyatt GH, Rennie D. Users’ guides to the medical literature [Editorial]. *JAMA*. 1993;270:2096-7. [PMID: 8411578]
37. Dubin A, Pozo MO, Casabella CA, Murias G, Pálizas F Jr, Moseinco MC, et al. Comparison of 6% hydroxyethyl starch 130/0.4 and saline solution for resuscitation of the microcirculation during the early goal-directed therapy of septic patients. *J Crit Care*. 2010;25:659.e1-8. [PMID: 20813485] doi:10.1016/j.jcrc.2010.04.007
38. Haupt MT, Rackow EC. Colloid osmotic pressure and fluid resuscitation with hetastarch, albumin, and saline solutions. *Crit Care Med*. 1982;10:159-62. [PMID: 6174274]
39. Li F, Sun H, Han XD. [The effect of different fluids on early fluid resuscitation in septic shock]. *Zhongguo Wei Zhong Bing Ji Jiu Yi Xue*. 2008;20:472-5. [PMID: 18687174]
40. Lv J, Zhao HY, Liu F, An YZ. [The influence of lactate Ringer solution versus hydroxyethyl starch on coagulation and fibrinolytic system in patients with septic shock]. *Zhongguo Wei Zhong Bing Ji Jiu Yi Xue*. 2012;24:38-41. [PMID: 22248750]
41. McIntyre LA, Fergusson D, Cook DJ, Rankin N, Dhingra V, Granton J, et al; Canadian Critical Care Trials Group. Fluid resuscitation in the management of early septic shock (FINESS): a randomized controlled feasibility trial. *Can J Anaesth*. 2008;55:819-26. [PMID: 19050085] doi:10.1007/BF03034053
42. Rackow EC, Mecher C, Astiz ME, Griffel M, Falk JL, Weil MH. Effects of pentastarch and albumin infusion on cardiorespiratory function and coagulation in patients with severe sepsis and systemic hypoperfusion. *Crit Care Med*. 1989;17:394-8. [PMID: 2468447]
43. Schortgen F, Lacherade JC, Bruneel F, Cattaneo I, Hemery F, Lemaire F, et al. Effects of hydroxyethylstarch and gelatin on renal function in severe sepsis: a multicentre randomised study. *Lancet*. 2001;357:911-6. [PMID: 11289347]
44. BaSES trial: Basel Starch Evaluation in Sepsis. *ClinicalTrials.gov*: NCT00273728. Accessed at <http://clinicaltrials.gov/ct2/show/record/NCT00273728> on 8 July 2014.
45. Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D, et al; SCCM/ESICM/ACCP/ATS/SIS. 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit Care Med*. 2003;31:1250-6. [PMID: 12682500]
46. Caironi P, Tognoni G, Masson S, Fumagalli R, Pesenti A, Romero M, et al; ALBIOS Study Investigators. Albumin replacement in patients with severe sepsis or septic shock. *N Engl J Med*. 2014;370:1412-21. [PMID: 24635772] doi:10.1056/NEJMoa1305727
47. Finfer S, McEvoy S, Bellomo R, McArthur C, Myburgh J, Norton R; SAFE Study Investigators. Impact of albumin compared to saline on organ function and mortality of patients with severe sepsis. *Intensive Care Med*. 2011;37:86-96. [PMID: 20924555] doi:10.1007/s00134-010-2039-6

Current Author Addresses: Dr. Rochweg: Critical Care Program Office, 2V3, McMaster University Medical Centre, 1200 Main Street West, Hamilton, Ontario L8N 3Z5, Canada.

Dr. Alhazzani: St. Joseph's Healthcare, 50 Charlton Avenue East, Hamilton, Ontario L8N 4A6, Canada.

Dr. Sindi: Department of Anesthesia and Critical Care, King Abdulaziz University Hospital, PO Box 80215, Jeddah 21589, Kingdom of Saudi Arabia.

Ms. Heels-Ansdell: Department of Clinical Epidemiology and Biostatistics, McMaster University, 1280 Main Street West, HSC-2C13, Hamilton, Ontario L8S 4K1, Canada.

Dr. Thabane: St. Joseph's Healthcare Hamilton, 3rd Floor, Martha Wing, Room H-325, 50 Charlton Avenue East, Hamilton, Ontario L8N 4A6, Canada.

Dr. Fox-Robichaud: McMaster University, DBRI C5-106, 237 Barton Street East, Hamilton, Ontario L2L 2X2, Canada.

Mr. Mbuagbaw: Department of Clinical Epidemiology and Biostatistics, Faculty of Health Sciences, McMaster University, 1280 Main Street West, HSC 2C7, Hamilton, Ontario L8S 4K1, Canada.

Drs. Szczeklik and Włodarczyk: Jagiellonian University Medical College, Skawińska 8, 31-066, Krakow, Poland.

Dr. Alshamsi: 1461 Upper Gage Avenue, Unit 7, Hamilton, Ontario L8W 1E6, Canada.

Dr. Altayyar: 67 Caroline Street South, Apartment 4A, Hamilton, Ontario L8P 3K6, Canada.

Dr. Ip: 602-10 John Street, Dundas, Ontario L9H 6J3, Canada.

Mr. Li: 49 Emerson Street, Hamilton, Ontario L8S 2X5, Canada.

Dr. Wang: Department of Internal Medicine, Hamilton General Hospital, 237 Barton Street East, Hamilton, Ontario L8L 3Z5, Canada.

Dr. Guyatt: McMaster University Health Sciences Centre, 1280 Main Street West, Room HSC-2C12, Hamilton, Ontario L8S 4K1, Canada.

Dr. Cook: Academic Chair, Critical Care Medicine, Room D176, St. Joseph's Healthcare, 50 Charlton Avenue East, Hamilton, Ontario L8N 3A5, Canada.

Dr. Jaeschke: 301 James Street South, Room F506, Hamilton, Ontario L8P 3B6, Canada.

Dr. Annane: General Intensive Care Unit, Raymond Poincaré Hospital (Assistance Publique-Hôpitaux de Paris), University of Versailles Saint-Quentin-en-Yvelines, 104 Boulevard Raymond Poincaré, 92380 Garches, France.

Author Contributions: Conception and design: B. Rochweg, W. Alhazzani, A. Sindi, L. Thabane, L. Mbuagbaw, W. Szczeklik, G.H. Guyatt, R. Jaeschke, D. Annane.

Analysis and interpretation of the data: B. Rochweg, W. Alhazzani, A. Sindi, D. Heels-Ansdell, L. Thabane, L. Mbuagbaw, W. Szczeklik, G.H. Guyatt, D.J. Cook, R. Jaeschke, D. Annane.

Drafting of the article: B. Rochweg, W. Alhazzani, A. Sindi, W. Szczeklik, R. Jaeschke, D. Annane.

Critical revision of the article for important intellectual content: B. Rochweg, W. Alhazzani, A. Sindi, L. Thabane, A. Fox-Robichaud, L. Mbuagbaw, W. Szczeklik, G.H. Guyatt, D.J. Cook, R. Jaeschke, D. Annane.

Final approval of the article: B. Rochweg, W. Alhazzani, A. Sindi, D. Heels-Ansdell, L. Thabane, A. Fox-Robichaud, L. Mbuagbaw, W. Szczeklik, G. Li, G.H. Guyatt, D.J. Cook, R. Jaeschke, D. Annane.

Provision of study materials or patients: D. Annane.

Statistical expertise: B. Rochweg, D. Heels-Ansdell, L. Thabane, L. Mbuagbaw, Q. Zhou.

Obtaining of funding: R. Jaeschke.

Administrative, technical, or logistic support: B. Rochweg, W. Alhazzani, A. Sindi.


Collection and assembly of data: B. Rochweg, W. Alhazzani, A. Sindi, L. Mbuagbaw, W. Szczeklik, F. Alshamsi, W.C. Ip, G. Li, A. Włodarczyk, R. Jaeschke, D. Annane.

Appendix Table. Risk of Bias, by Study

Study, Year (Reference)	Randomization-Sequence Generation	Randomization Concealment	Blinding	Incomplete Data	Selective Reporting	Other*	Overall for Mortality
Myburgh et al, 2012 (3)	Low	Low	Low	Low	Low	Low	Low
Guidet et al, 2012 (6)	Low	Low	Low	Low	Low	Low	Low
Perner et al, 2012 (4)	Low	Low	Low	Low	Low	Low	Low
Haupt and Rackow, 1982 (38)	Probably low	Probably low	Low	Low	Low	Low	Probably low
Rackow et al, 1989 (42)	Probably low	Probably low	Probably low	Low	Low	Low	Probably low
Brunkhorst et al, 2008 (1)	Low	Low	Low	Low	Low	Low	Low
Dubin et al, 2010 (37)	Low	Probably low	Low	Low	Low	Low	Probably low
Li et al, 2008 (39)	Probably low	Probably low	Probably low	Probably low	Probably low	Probably low	Probably low
Lv et al, 2012 (40)	Probably low	Probably low	Probably low	Probably low	Probably low	Probably low	Probably low
McIntyre et al, 2008 (41)	Low	Low	Low	Low	Low	Low	Low
Finfer et al, 2004 (2)	Low	Low	Low	Low	Low	Low	Low
Annane et al, 2013 (16)	Low	Probably low	Low	Low	Low	Low	Probably low
Siegemund, 2011 (44)	Low	Low	Low	Low	Low	Low	Low
Schortgen et al, 2001 (43)	Low	Low	Low	Low	Low	Low	Low


* Including intention-to-treat analysis.

Appendix Figure 1. Network map for 4-node analysis.




HES = hydroxyethyl starch.

Appendix Figure 2. Network map for 6-node analysis.



BC = balanced crystalloid; H-HES = high-molecular-weight hydroxyethyl starch; L-HES = low-molecular-weight hydroxyethyl starch.

Appendix Figure 3. Forest plot for mortality in direct comparisons of all crystalloids vs. all colloids.



M-H = Mantel-Haenszel.